



ОИЯИ
ОБЪЕДИНЁННЫЙ ИНСТИТУТ
ЯДЕРНЫХ ИССЛЕДОВАНИЙ



Старая Стекловка, R-операция и ренормгруппа

Д.Казаков

Лаборатория теоретической физики им. Н.Н.Боголюбова

Объединённый институт ядерных исследований

13 ноября '25

Дубна

Renormalization

Bogoliubov-Parasiuk-Hepp-Zimmermann R-operation

$$RG = \prod_{\gamma_i} (1 - M_i) G$$

$$RG = (1 - K) R' G$$

G - graph

γ - divergent subgraph

M_i - subtraction operator

K-operation extracts the singular part

Incomplete R-operation

A.N.Vasiliev, Green Book


$$R' = 1 - \sum_{\gamma} K R' \gamma + \sum_{\gamma, \gamma'} K R' \gamma K R' \gamma' - \dots;$$

R-operation is equivalent to the introduction of the counterterms into the Lagrangian

$$\mathcal{L} \Rightarrow \mathcal{L} + \Delta \mathcal{L}$$

Bogolyubov-Parasiuk Theorem: In any local quantum field theory after subtracting the UV divergences in subgraphs the resulting counterterms are always local in coordinate space or at most are polynomials of external momenta in momentum space in each order of perturbation theory

BPHZ R-operation



- dimensional regularization
 $d = 4 - 2\epsilon$

$$\begin{aligned} \mathcal{R}' G_n = & \frac{A_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^n} + \frac{A_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^n} + \dots + \frac{A_1^{(n)} (\mu^2)^\epsilon}{\epsilon^n} \\ & + \frac{B_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^{n-1}} + \frac{B_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^{n-1}} + \dots + \frac{B_1^{(n)} (\mu^2)^\epsilon}{\epsilon^{n-1}} \\ & + \frac{C_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^{n-2}} + \frac{C_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^{n-2}} + \dots + \frac{C_1^{(n)} (\mu^2)^\epsilon}{\epsilon^{n-2}} \\ & + \text{lower pole terms,} \end{aligned}$$

$A_k^{(n)} \quad B_k^{(n)} \quad C_k^{(n)} \quad (\mu^2)^{k\epsilon}$ terms appear after subtraction of (n-k) loop counter terms

BPHZ R-operation

Bogoliubov-Parasiuk Theorem: (Locality)

$R'G_n$ is local, i.e. terms like $\log^k \mu^2 / \epsilon^m$ should cancel for any k and m

- Due to locality all higher order divergences are related to the lower ones

$$A_n^{(n)} = (-1)^{n+1} \frac{A_1^{(n)}}{n}, \quad \text{One loop}$$

$$B_n^{(n)} = (-1)^n \left(\frac{2}{n} B_2^{(n)} + \frac{n-2}{n} B_1^{(n)} \right), \quad \text{One and two loops}$$

$$C_n^{(n)} = (-1)^{n+1} \left(\frac{3}{n} C_3^{(n)} + \frac{2(n-3)}{n} C_2^{(n)} + \frac{(n-2)(n-3)}{2n} C_1^{(n)} \right). \quad \text{One, two and three loops}$$

$$\mathcal{KR}'G_n = \sum_{k=1}^n \left(\frac{A_k^{(n)}}{\epsilon^n} + \frac{B_k^{(n)}}{\epsilon^{n-1}} + \frac{C_k^{(n)}}{\epsilon^{n-2}} + \dots \right) \equiv \frac{A_n^{(n)'}}{\epsilon^n} + \frac{B_n^{(n)'}}{\epsilon^{n-1}} + \frac{C_n^{(n)'}}{\epsilon^{n-2}} + \dots$$

$$A_n^{(n)'} = (-1)^{n+1} A_n^{(n)} = \frac{A_1^{(n)}}{n}, \quad \text{One loop}$$

$$B_n^{(n)'} = \left(\frac{2}{n(n-1)} B_2^{(n)} + \frac{2}{n} B_1^{(n)} \right), \quad \text{One and two loops}$$

$$C_n^{(n)'} = \left(\frac{2}{(n-1)(n-2)} \frac{3}{n} C_3^{(n)} + \frac{2}{n-1} \frac{3}{n} C_2^{(n)} + \frac{3}{n} C_1^{(n)} \right). \quad \text{One, two and three loops}$$

The Recurrence Relations

- These properties allow one to write down the recurrence relations connecting the subsequent orders of the counterterms and to evaluate them algebraically without calculating the diagrams. This can be done in renormalizable and non-renormalizable theories. The difference is a more complicated structure of these relations in NR case.

Leading divergences:

$$\begin{aligned}
 n \quad & \text{[Diagram: a solid circle]} = - \text{[Diagram: a dashed circle with a solid circle inside and a tadpole on the left]} - \text{[Diagram: a dashed circle with a solid circle inside and a tadpole on the right]} + \sum_{k=1}^{n-2} \text{[Diagram: a dashed circle with a solid circle inside and a tadpole on the left]} \text{[Diagram: a dashed circle with a solid circle inside and a tadpole on the right]} \\
 \text{\textit{n-loop}} \quad & \text{\textit{(n)}} \quad \text{\textit{(n-1)-loop}} \quad \text{\textit{(n-1)-loop}} \quad \text{\textit{k-loop}} \quad \text{\textit{(n-k-1)-loop}} \\
 A_n^{(n)} \quad & A_{n-1}^{(n)} \quad A_{n-1}^{(n)} \quad A_k^{(n)} \quad A_{n-k-1}^{(n)}
 \end{aligned}$$

- These recurrence relations can be promoted to the RG equations for the scattering amplitudes, effective potential, etc which sum up the leading divergences (logarithms) and to find out the high energy/field behaviour

$$\phi_6^4$$

Loop Expansion (non-renormalizable case)

UV divergences
within dim reg

$$\Delta\mathcal{L}_1 \sim \lambda^2(s+t+u)\Phi^4\left(\frac{1}{\epsilon} + c_{11}\right)$$

$$\Delta\mathcal{L}_1 \sim \lambda^2\partial^2\Phi^2\Phi^2\left(\frac{1}{\epsilon} + c_{11}\right),$$

Momentum space

Coordinate space

Increasing power of momentum

$$\Delta\mathcal{L} = \lambda^2\partial^2\Phi^2\Phi^2\left(\frac{1}{\epsilon} + c_{11}\right) + \lambda^3\left[\partial^4\Phi^2\Phi^2\left(\frac{1}{\epsilon^2} + \frac{1}{\epsilon} + c_{12}\right) + \partial^2\Phi^2\partial^2\Phi^2\left(\frac{1}{\epsilon^2} + \frac{1}{\epsilon} + c_{13}\right)\right] + \lambda^4[\dots] + \lambda^5[\dots]$$

$$\lambda^3\Phi^6\left(\frac{1}{\epsilon} + c_{21}\right) + \lambda^4\left[\partial^2\Phi^4\Phi^2\left(\frac{1}{\epsilon^2} + \frac{1}{\epsilon} + c_{22}\right) + \partial^2\Phi^2\Phi^4\left(\frac{1}{\epsilon^2} + \frac{1}{\epsilon} + c_{23}\right)\right]$$

Increasing power of fields

$$\lambda^5\Phi^8\left(\frac{1}{\epsilon^2} + \frac{1}{\epsilon} + c_{32}\right),$$



D=6 N=2**S-channel** $S_n(s, t)$ **T-channel** $T_n(s, t)$ $T_n(s, t) = S_n(t, s)$ **Exact all-loop recurrence relation** $S_3 = -s/3, T_3 = -t/3$

$$nS_n(s, t) = -2s \int_0^1 dx \int_0^x dy (S_{n-1}(s, t') + T_{n-1}(s, t'))$$


 $n \geq 4$ $t' = t(x - y) - sy$ **D=8 N=1****S-channel** $S_n(s, t)$ **T-channel** $T_n(s, t)$ $T_n(s, t) = S_n(t, s)$ **Exact all-loop recurrence relation** $S_1 = \frac{1}{12}, T_1 = \frac{1}{12}$

$$\begin{aligned}
 nS_n(s, t) = & -2s^2 \int_0^1 dx \int_0^x dy y(1-x) (S_{n-1}(s, t') + T_{n-1}(s, t'))|_{t'=tx+yu} \\
 + & s^4 \int_0^1 dx x^2(1-x)^2 \sum_{k=1}^{n-2} \sum_{p=0}^{2k-2} \frac{1}{p!(p+2)!} \frac{d^p}{dt'^p} (S_k(s, t') + T_k(s, t')) \times \\
 & \times \frac{d^p}{dt'^p} (S_{n-1-k}(s, t') + T_{n-1-k}(s, t'))|_{t'=-sx} (tsx(1-x))^p
 \end{aligned}$$

Effective Potential in Scalar Theory

V_{eff} Is the sum of all vacuum IPI diagrams

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)^2 - gV_0(\phi)$$



$$v_2(\phi) \equiv \frac{d^2 V_0(\phi)}{d\phi^2}$$

$$v_n \equiv d^n V_0 / d\phi^n$$


Shown are UV divergent vacuum diagrams in arbitrary scalar theory up to three loops

$$V_{eff} = g \sum_{n=0}^{\infty} (-g)^n V_n.$$

Recurrence relations for the leading poles for effective potential

Kazakov, Iakhibbaev, Tolkachev 22

Action of R' -operation on divergent diagram




$$D_2 = \frac{d^2}{d\phi^2}$$

$$n\Delta V_n = \frac{1}{2}v_2 D_2\Delta V_{n-1} + \frac{1}{4} \sum_{k=1}^{n-2} D_2\Delta V_k D_2\Delta V_{n-1-k}, \quad n \geq 2 \quad \Delta V_1 = \frac{1}{4}v_2^2$$

$$n\Delta V_n = \frac{1}{4} \sum_{k=0}^{n-1} D_2\Delta V_k D_2\Delta V_{n-1-k}, \quad n \geq 1, \quad \Delta V_0 = V_0$$

From Recurrence Relation to the RG Equation



$$n \text{ } \underbrace{\text{solid circle}}_{\substack{n\text{-loop} \\ A_n^{(n)}}} = - \underbrace{\text{solid circle} \text{ --- } \text{dashed circle}}_{\substack{(n-1)\text{-loop} \\ A_{n-1}^{(n)}}} - \underbrace{\text{dashed circle} \text{ --- } \text{solid circle}}_{\substack{(n-1)\text{-loop} \\ A_{n-1}^{(n)}}} + \sum_{k=1}^{n-2} \underbrace{\text{solid circle} \text{ --- } \text{dashed circle}}_{\substack{k\text{-loop} \\ A_k^{(n)}}} \underbrace{\text{solid circle}}_{\substack{(n-k-1)\text{-loop} \\ A_{n-k-1}^{(n)}}}$$

- This is the general recurrence relation that reflects the locality of counterterms in any theory
- In renormalizable theories A_n is a constant and this relation is reduced to the algebraic one
- In non-renormalizable theories for the scattering amplitudes A_n depends on kinematics and one has to integrate through the one-loop diagrams
- For the effective potential A_n depends on the fields and one has to differentiate

Taking the sum $\sum_n A_n (-z)^n = A(z)$ one can transform the recurrence relation into RG equation

$$\frac{dA(z)}{dz} = -2 \oint A(z) + \oint A(z) \otimes A(z)$$

$$\frac{d}{dz} = \frac{d}{d \log \mu^2}$$

This is the generalized RG equation valid in any (even non-renormalizable) theory!

RG Equation

SYM_D

D=6 N=2

$$\Sigma(s, t, z) = z^{-2} \sum_{n=3}^{\infty} (-z)^n S_n(s, t)$$

$$\frac{d}{dz} \Sigma(s, t, z) = s - \frac{2}{z} \Sigma(s, t, z) + 2s \int_0^1 dx \int_0^x dy (\Sigma(s, t', z) + \Sigma(t', s, z))|_{t'=xt+yu}$$

Linear equation

D=8 N=1

$$\Sigma(s, t, z) = \sum_{n=1}^{\infty} (-z)^n S_n(s, t)$$

$$\begin{aligned} \frac{d}{dz} \Sigma(s, t, z) = & -\frac{1}{12} + 2s^2 \int_0^1 dx \int_0^x dy y(1-x) (\Sigma(s, t', z) + \Sigma(t', s, z))|_{t'=tx+yu} \\ & -s^4 \int_0^1 dx x^2(1-x)^2 \sum_{p=0}^{\infty} \frac{1}{p!(p+2)!} \left(\frac{d^p}{dt'^p} (\Sigma(s, t', z) + \Sigma(t', s, z))|_{t'=-sx} \right)^2 (tsx(1-x))^p. \end{aligned}$$

Non-linear equation

RG pole equation for arbitrary potential

$$\Sigma(z, \phi) = \sum_{n=0}^{\infty} (-z)^n \Delta V_n(\phi) \quad z = \frac{g}{\epsilon}$$

Kazakov, Iakhibbaev, Tolkachev 22

JHEP v 2304 (2023) 128

RG pole equation

$$\frac{d\Sigma}{dz} = -\frac{1}{4} (D_2 \Sigma)^2 \quad \Sigma(0, \phi) = V_0(\phi)$$

This a non-linear partial differential equation!

Effective potential

$$V_{eff}(g, \phi) = g \Sigma(z, \phi) \Big|_{z \rightarrow -\frac{g}{16\pi^2} \log g v_2 / \mu^2} \quad v_2(\phi) \equiv \frac{d^2 V_0(\phi)}{d\phi^2}$$

RG Equations in Subleading Order Scattering Amplitude

$$\frac{dS(s, t, z)}{dz} = s(S_1 + T_1) \otimes (S_1 + T_1) \quad \text{Coefficients of } 1/\epsilon^n$$

$$\begin{aligned} \frac{d^2 S_2(s, t, z)}{dz^2} = & s \frac{d}{dz} (S_1 + T_1) \otimes (S_2 + T_2) \\ & + s^2 (S_1 + T_1) \otimes (S_2 + T_2) \otimes (S_1 + T_1) \\ & + s (S_1 + T_1) \otimes \otimes (S_1 + T_1) \end{aligned} \quad \begin{array}{l} \text{Coefficients of} \\ 1/\epsilon^{n-1} \end{array}$$

- ◆ Equation for the subleading order function as well as for all the subsequent orders is always linear!
- ◆ This seems to be in contradiction with the usual RG equation but is not!

RG Equations in Subleading Order

Effective potential

$$\frac{\partial \Sigma_A}{\partial z} = -\frac{1}{4}(D_2 \Sigma_A)^2$$

Σ_A - leading order

Σ_B - subleading order

Σ_G - kinetic term

$$\begin{aligned} \frac{\partial^2 \Sigma_B}{\partial z^2} = & -2 \frac{1}{4} \frac{\partial}{\partial z} \left\{ 2 D_2 \Sigma_A D_2 \Sigma_B - (D_2 \Sigma_A)^2 D_2 \Sigma_G \right\} - \\ & -2 \left\{ \frac{1}{8} (D_3 \Sigma_A)^2 D_2 \Sigma_B - 2 \frac{1}{8} (D_3 \Sigma_A)^2 D_2 \Sigma_A D_2 \Sigma_B + 2 \frac{1}{8} D_3 \Sigma_A D_3 \Sigma_B D_2 \Sigma_A + \right. \\ & + 2 \frac{1}{8} D_3 \Sigma_A D_3 \Sigma_B D_2 \Sigma_A - \frac{1}{8} (D_3 \Sigma_A)^2 D_2 \Sigma_A + \frac{1}{8} (D_2 \Sigma_A)^2 D_4 \Sigma_B - \\ & \left. - 2 \frac{1}{8} (D_2 \Sigma_A)^2 D_4 \Sigma_A D_2 \Sigma_G + 2 \frac{1}{8} D_2 \Sigma_A D_2 \Sigma_B D_4 \Sigma_A \right\} \end{aligned}$$

$$D_n \equiv \frac{\partial^n}{\partial \phi^n}$$

$$\frac{\partial^2 \Sigma_G}{\partial z^2} = \frac{1}{12} (D_3 \Sigma_A)^2$$

- ◆ Equation for the subleading order function as well as for all the subsequent orders is always linear!
- ◆ This seems to be in contradiction with the usual RG equation which is non-linear, however, the proper form of the usual RG equation is also linear!!

V-A four fermion Interaction

Lagrangian

$$\mathcal{L} = i\bar{\Psi}\hat{\partial}\Psi - \frac{G}{\sqrt{2}}(\bar{\Psi}\hat{\mathcal{O}}\Psi)(\bar{\Psi}\hat{\mathcal{O}}\Psi) \quad \hat{\mathcal{O}} = \gamma^\mu(1 - \gamma^5)/2$$

Tree level amplitude

$$A_4^{(0)} = \langle 13 \rangle [42]$$

One-loop amplitude

$$A_4^{(1)} = S_1 + T_1 + U_1 = \left(-\frac{16}{3} \frac{s}{\epsilon} - \frac{16}{3} \frac{t}{\epsilon} \right) \langle 13 \rangle [42]$$

RG Equation for the Scattering Amplitude

$$A(s, t, u) = \sum_{n=0}^{\infty} A_n(s, t, u)(-z)^n$$

$$\begin{aligned} -\frac{dA(s, t, u)}{dz} = & \\ & -4s \int_0^1 dx \sum_{p=0}^{\infty} \frac{[s(-s-u)]^p [x(1-x)]^{p+1}}{p!(p+1)!(p+2)^{-1}} \left(\frac{d^p A(s, -s-u', u')}{du'^p} \right)^2 \\ & -4t \int_0^1 dx \sum_{p=0}^{\infty} \frac{[t(-t-u)]^p [x(1-x)]^{p+1}}{p!(p+1)!(p+2)^{-1}} \left(\frac{d^p A(-t-u', t, u')}{du'^p} \right)^2 \\ & +4u \int_0^1 dx \sum_{p=0}^{\infty} \frac{[u(-s-u)]^p [x(1-x)]^{p+1}}{p!(p+1)!(p+3)^{-1}} \left(\frac{d^p A(s', -s'-u, u)}{ds'^p} \right)^2 \\ & +4u \int_0^1 dx \sum_{p=0}^{\infty} \frac{[u(-t-u)]^p [x(1-x)]^{p+1}}{p!(p+1)!(p+3)^{-1}} \left(\frac{d^p A(-t'-u, t', u)}{dt'^p} \right)^2 \end{aligned}$$


$$z = G/\epsilon \rightarrow -G \log Q^2/\mu^2$$

- ◆ With the help of this equation one can find the asymptotic behaviour of the amplitude in the high energy regime $s \sim t \sim u \sim E^2 \rightarrow \infty$

Numerical Solution of RG Equation for the Scattering Amplitude

Borlakov, Kazakov, 25

$$s = 4E^2, \quad t = u = -2E^2, \quad E \rightarrow \infty \quad \frac{\text{Amplitude}}{A_{Tree}} = A(E^2, G \log E^2 / \mu^2)$$



$$A_{Tree} \sim GE^2 \quad A \sim \frac{1}{GE^2 \log E^2} \quad Amp = A_{Tree} A \sim \frac{1}{\log E^2}$$

α -attractor Inflaton Potential

Inflaton action with hyperbolic geometry

Kallosh, Linde 13

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R(g) + \frac{1}{2} \frac{\partial_\mu \phi \partial^\mu \phi}{1 - \frac{\phi^2}{6\alpha}} - V(\phi) \right]$$

Transition to the standard kinetic term

$$\partial\phi / \sqrt{1 - \frac{\phi^2}{6\alpha}} = \partial\varphi \quad \phi = \sqrt{6\alpha} \tanh\left(\frac{\varphi}{\sqrt{6\alpha}}\right)$$

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R(g) + \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - V\left(\sqrt{6\alpha} \tanh\left(\frac{\varphi}{\sqrt{6\alpha}}\right)\right) \right].$$

T_2 - model

$$V(\phi) = g\phi^2$$



$$V_T(\varphi) = \tanh^2\left(\frac{\varphi}{\sqrt{6\alpha}M_{Pl}}\right)$$

$$\frac{d\Sigma}{dz} = -\frac{1}{4}(D_2\Sigma)^2$$

Dimensionless variables

$$x = z/M_{Pl}^4 \quad y = \tanh^n(\varphi/\sqrt{6\alpha}M_{Pl})$$

$$\Sigma(z/M_{Pl}^4, \tanh^n(\varphi/\sqrt{6\alpha}M_{Pl})) \equiv S(x, y)$$

$$S_x = -\frac{(y-1)^2 ((3y-1)S_y + 2(y-1)yS_{yy})^2}{36\alpha^2}$$

$$S(0, y) = y, \quad S(x, 1) = 1, \quad S_y(x, 1) = 0.$$


Lift of the Potential at the Minima - Origin of the Cosmological Constant



Kazakov, Iakhibbaev, Tolkachev, Filippov 24,25

ArXiv: 2405.18818

Comparison of the classical T2-model potential (blue dashed line), the one-loop correction (orange dashed line), and the RG summed potential (black solid line) for $g \sim 1, \mu < M_{Pl}$




T2-model potential: variation of μ . The classical potential (blue dashed line), the RG summed potential (solid lines) for

$$\mu < M_{Pl} \quad \mu \ll M_{Pl} \quad \mu > M_{Pl}$$

black line, red line, green line

$$g = 2, \alpha = 1$$



Resume

- **The UV divergences in non-renormalizable theories are local and can be removed by local counter terms like in renormalizable ones**
- **Based on locality of the counter terms due to the Bogoliubov-Parasiuk theorem one can construct the recurrence relations for the leading, subleading, etc divergences in all loops divergences starting from one-, two-, three-, etc loop diagrams**
- **The recurrence relations can be converted into the generalized RG equations just like in renormalizable theories**
- **The RG equations allow one to sum up the leading (subleading, etc) divergences in all loops and define the high-energy/field behaviour**
- **The RG equations for the subleading poles, etc are always linear differential equation of increasing order**
- **The RG equations for logarithms are always first order differential equation with known rhs**